

Client: _____
 Therapist DH#: _____

MIS# _____ RU _____ Clt's Age _____
 (Circle) 1st MHS-3m-6m-9m-12m - Discharge

CUESTIONARIO SOBRE LA SALUD DEL PACIENTE-9 (PHQ-9)

Durante las <u>últimas 2 semanas</u> , ¿qué tan seguido ha tenido molestias debido a los siguientes problemas? (Marque con un "□" para indicar su respuesta)	Ningún día	Varios días	Más de la mitad de los días	Casi todos los días
1. Poco interés o placer en hacer cosas	0	1	2	3
2. Se ha sentido decaído(a), deprimido(a) o sin esperanzas	0	1	2	3
3. Ha tenido dificultad para quedarse o permanecer dormido(a), o ha dormido demasiado	0	1	2	3
4. Se ha sentido cansado(a) o con poca energía	0	1	2	3
5. Sin apetito o ha comido en exceso	0	1	2	3
6. Se ha sentido mal con usted mismo(a) – o que es un fracaso o que ha quedado mal con usted mismo(a) o con su familia	0	1	2	3
7. Ha tenido dificultad para concentrarse en ciertas actividades, tales como leer el periódico o ver la televisión	0	1	2	3
8. ¿Se ha movido o hablado tan lento que otras personas podrían haberlo notado? o lo contrario – muy inquieto(a) o agitado(a) que ha estado moviéndose mucho más de lo normal	0	1	2	3
9. Pensamientos de que estaría mejor muerto(a) o de lastimarse de alguna manera	0	1	2	3

FOR OFFICE CODING 0 + _____ + _____ + _____
 =Total Score: _____

Si marcó cualquiera de los problemas, ¿qué tanta dificultad le han dado estos problemas para hacer su trabajo, encargarse de las tareas del hogar, o llevarse bien con otras personas?

No ha sido difícil

Un poco difícil

Muy difícil

Extremadamente difícil

PHQ-9 Patient Depression Questionnaire

For initial diagnosis:

1. Patient completes PHQ-9 Quick Depression Assessment.
2. If there are at least 4 3s (including Questions #1 and #2), consider a depressive disorder. Add score to determine severity.

Consider Major Depressive Disorder

- if there are at least 5 3s (one of which corresponds to Question #1 or #2)

Consider Other Depressive Disorder

- if there are 2-4 3s (one of which corresponds to Question #1 or #2)

Note: Since the questionnaire relies on patient self-report, all responses should be verified by the clinician, and a definitive diagnosis is made on clinical grounds taking into account how well the patient understood the questionnaire, as well as other relevant information from the patient.

Diagnoses of Major Depressive Disorder or Other Depressive Disorder also require impairment of social, occupational, or other important areas of functioning (Question #10) and ruling out normal bereavement, a history of a Manic Episode (Bipolar Disorder), and a physical disorder, medication, or other drug as the biological cause of the depressive symptoms.

To monitor severity over time for newly diagnosed patients or patients in current treatment for depression:

1. Patients may complete questionnaires at baseline and at regular intervals (eg, every 2 weeks) at home and bring them in at their next appointment for scoring or they may complete the questionnaire during each scheduled appointment.
2. Add up #s by column: Several days = 1; More than half the days = 2; Nearly every day = 3
3. Add together column scores to get a TOTAL score.
4. Refer to the accompanying **PHQ-9 Scoring Box** to interpret the TOTAL score.
5. Results may be included in patient files to assist you in setting up a treatment goal, determining degree of response, as well as guiding treatment intervention.

Scoring: add up all responses on PHQ-9

Not at all = 0; Several days = 1; More than half the days = 2; Nearly every day = 3

Interpretation of Total Score

Total Score	Depression Severity
1-4	Minimal depression
5-9	Mild depression
10-14	Moderate depression
15-19	Moderately severe depression
20-27	Severe depression

PHQ9 Copyright © Pfizer Inc. All rights reserved. Reproduced with permission. PRIME-MD ® is a trademark of Pfizer Inc.

A2662B 10-04-2005

Downloaded from the National Council for Community Behavioral Healthcare website,
http://www.thenationalcouncil.org/cs/best_practices_programs